

Online-Only Abstracts

Dissemination of New Delhi metallo- β -lactamase-I-producing *Acinetobacter baumannii* in Europe

R. A. Bonnin¹, L. Poirel¹, T. Naas¹, M. Pirs², K. Seme², J. Schrenzel³ and P. Nordmann¹

1) Service de Bactériologie-Virologie, INSERM U914 “Emerging Resistance to Antibiotics”, Hôpital de Bicêtre, Assistance Publique/Hôpitaux de Paris, Faculté de Médecine et Université Paris-Sud, K-Bicêtre, France, 2) Institute of Microbiology and Immunology Medical Faculty, University of Ljubljana, Ljubljana, Slovenia and 3) Bacteriology Laboratory and Genomic Research Laboratory, Geneva University Hospitals, Geneva, Switzerland

Original Submission: 15 February 2012; **Revised Submission:** 20 April 2012; **Accepted:** 20 May 2012

Editor: R. Cantón

Article published online: 25 May 2012

Clin Microbiol Infect 2012; **18**: E362–E365

10.1111/j.1469-0691.2012.03928.x

Abstract

Multidrug-resistant and New Delhi metallo- β -lactamase I (NDM-I) -producing *Acinetobacter baumannii* are increasingly reported. A collection of five NDM-I-positive *A. baumannii* isolates recovered in four European countries were analysed. Genotyping was performed by pulsed-field gel electrophoresis, multiplex PCR sequence typing, Diversilab and multilocus sequence typing. Three distinct sequence types were identified. All isolates harboured a chromosomally located *bla*_{NDM-I} gene within a Tn125-like transposon. One isolate co-expressed another unrelated carbapenemase OXA-23. This report constitutes the first epidemiological study of NDM-I-producing *A. baumannii* from four countries.

Multidrug resistance-encoding plasmid from *Aeromonas* sp. strain P2G1

E. Marti and J. L. Balcázar

Catalan Institute for Water Research (ICRA), Scientific and Technological Park of the University of Girona, Girona, Spain

Original Submission: 21 February 2012; **Revised Submission:** 22 May 2012; **Accepted:** 25 May 2012

Editor: J.-M. Rolain

Article published online: 30 May 2012

Clin Microbiol Infect 2012; **18**: E366–E368

10.1111/j.1469-0691.2012.03935.x

Abstract

A plasmid (pP2G1), which confers multidrug resistance in an environmental *Aeromonas* species, was completely sequenced using a shotgun approach. Plasmid pP2G1 encoded resistance to aminoglycosides and quinolones [*aac*(6′)-Ib-cr], β -lactams (*bla*_{OXA-1}), chloramphenicol (*catB3*), macrolides [*mphA-mrx-mphR*], quaternary ammonium compounds (*qacEΔ1*), quinolones (*qnrS2*), rifampicin (*arr-3*) and sulphonamides (*sul1*). These findings suggest that *Aeromonas* species may potentially act as reservoirs of antibiotic resistance genes.

Metallo- β -lactamase-producing *Pseudomonas aeruginosa* in the Netherlands: the nationwide emergence of a single sequence type

A. K. Van der Bijl^{1,2}, D. Van der Zwan¹, G. Peirano³, J. A. Severin¹, J. D. D. Pitout³, M. Van Westreenen¹, W. H. F. Goessens¹, on behalf of the MBL-PA surveillance study group

1) Department of Medical Microbiology and Infectious Diseases, Erasmus University Medical Centre, Rotterdam, 2) Department of Microbiology, Reinier de Graaf Groep, Delft, the Netherlands and 3) Department of Pathology and Laboratory Medicine, Microbiology and Infectious Diseases, University of Calgary, Calgary, AB, Canada

Original Submission: 5 March 2012; **Revised Submission:** 18 May 2012; **Accepted:** 10 June 2012

Editor: R. Cantón

Article published online: 14 June 2012

Clin Microbiol Infect 2012; **18**: E369–E372

10.1111/j.1469-0691.2012.03969.x

Abstract

Recently, the first outbreak of clonally related VIM-2 metallo- β -lactamase (MBL)-producing *Pseudomonas aeruginosa* in a Dutch tertiary-care centre was described. Subsequently, a nationwide surveillance study was performed in 2010–2011, which identified the presence of VIM-2 MBL-producing *P. aeruginosa* in 11 different hospitals. Genotyping by multiple-locus variable-number tandem-repeat analysis (MLVA) showed that the majority of the 82 MBL-producing isolates found belonged to a single MLVA type ($n = 70$, 85%), identified as ST111 by multilocus sequence typing (MLST). As MBL-producing isolates cause serious infections that are difficult to treat, the presence of clonally related isolates in various hospitals throughout the Netherlands is of nationwide concern.